ORIGINAL ARTICLE

A Trial of Intraoperative Low-Tidal-Volume Ventilation in Abdominal Surgery

Emmanuel Futier, M.D., Jean-Michel Constantin, M.D., Ph.D., Catherine Paugam-Burtz, M.D., Ph.D., Julien Pascal, M.D., Mathilde Eurin, M.D., Arthur Neuschwander, M.D., Emmanuel Marret, M.D., Marc Beaussier, M.D., Ph.D., Christophe Gutton, M.D., Jean-Yves Lefrant, M.D., Ph.D., Bernard Allaouchiche, M.D., Ph.D., Daniel Verzilli, M.D., Marc Leone, M.D., Ph.D., Audrey De Jong, M.D., Jean-Etienne Bazin, M.D., Ph.D., Bruno Pereira, Ph.D., and Samir Jaber, M.D., Ph.D., for the IMPROVE Study Group*

ABSTRACT

BACKGROUND

Lung-protective ventilation with the use of low tidal volumes and positive end-expiratory pressure is considered best practice in the care of many critically ill patients. However, its role in anesthetized patients undergoing major surgery is not known.

METHODS

In this multicenter, double-blind, parallel-group trial, we randomly assigned 400 adults at intermediate to high risk of pulmonary complications after major abdominal surgery to either nonprotective mechanical ventilation or a strategy of lung-protective ventilation. The primary outcome was a composite of major pulmonary and extrapulmonary complications occurring within the first 7 days after surgery.

RESULTS

The two intervention groups had similar characteristics at baseline. In the intention-to-treat analysis, the primary outcome occurred in 21 of 200 patients (10.5%) assigned to lung-protective ventilation, as compared with 55 of 200 (27.5%) assigned to nonprotective ventilation (relative risk, 0.40; 95% confidence interval [CI], 0.24 to 0.68; P=0.001). Over the 7-day postoperative period, 10 patients (5.0%) assigned to lung-protective ventilation required noninvasive ventilation or intubation for acute respiratory failure, as compared with 34 (17.0%) assigned to nonprotective ventilation (relative risk, 0.29; 95% CI, 0.14 to 0.61; P=0.001). The length of the hospital stay was shorter among patients receiving lung-protective ventilation than among those receiving nonprotective ventilation (mean difference, –2.45 days; 95% CI, –4.17 to –0.72; P=0.006).

CONCLUSIONS

As compared with a practice of nonprotective mechanical ventilation, the use of a lung-protective ventilation strategy in intermediate-risk and high-risk patients undergoing major abdominal surgery was associated with improved clinical outcomes and reduced health care utilization. (IMPROVE ClinicalTrials.gov number, NCT01282996.)

From the Département d'Anesthésie et Réanimation, Hôpital Estaing (E.F., J.-M.C., J.P., J.-E.B.), Université de Clermont-Ferrand, Retinoids, Reproduction, and Developmental Diseases Unit, Équipe Accueil 7281 (E.F., J.-M.C.), and the Biostatistics Unit, Direction de la Recherche Clinique (B.P.), Centre Hospitalier Universitaire (CHU) de Clermont-Ferrand, Clermont-Ferrand; Assistance Publique–Hôpitaux de Paris (AP-HP), Département d'Anesthésie et Réanimation, Hôpital Beaujon, Hôpitaux Universitaires Paris Nord Val de Seine and Université Paris Diderot, Sorbonne Paris Cité (C.P.-B., M.E., A.N.), Département d'Anesthésie et Réanimation, Hôpital Tenon (E.M.), and AP-HP, Département d'Anesthésie et Réanimation, Hôpital Saint-Antoine (M.B., C.G.), Paris; CHU de Nîmes, Section d'Anesthésie and Département d'Anesthésie et Réanimation, Nîmes (J.-Y.L.); CHU de Lyon, Département d'Anesthésie et Réanimation, Hôpital Edouard Herriot, Lyon (B.A.); CHU de Montpellier, Département d'Anesthésie et Réanimation B, Hôpital Saint-Eloi, and INSERM Unité 1046 and Université Montpellier 1, Montpellier (D.V., A.D.J., S.J.); and Assistance Publique–Hôpital de Marseille, Département d'Anesthésie et Réanimation, Hôpital Nord, Marseille (M.L.) — all in France. Address reprint requests to Dr. Jaber at the Département d'Anesthésie et Réanimation B (DAR B), 80 Ave. Augustin Fliche, 34295 Montpellier, France, or at s-jaber@chu-montpellier.fr.

*Additional investigators in the Intraoperative Protective Ventilation (IMPROVE) Study Group are listed in the Supplementary Appendix, available at NEJM.org.

N Engl J Med 2013;369:428-37.
DOI: 10.1056/NEJMoa1301082
Copyright © 2013 Massachusetts Medical Society.

The New England Journal of Medicine is produced by NEJM Group, a division of the Massachusetts Medical Society.
Downloaded from nejm.org on October 14, 2026.

Worldwide, more than 230 million patients undergoing major surgery each year require general anesthesia and mechanical ventilation.[1] Postoperative pulmonary complications adversely affect clinical outcomes and health care utilization,[2] so prevention of these complications has become a measure of the quality of hospital care.[3] Previous, large cohort studies have shown that 20 to 30% of patients undergoing surgery with general anesthesia are at intermediate to high risk for postoperative pulmonary complications.[4,5]

Mechanical ventilation with the use of high tidal volumes (10 to 15 ml per kilogram of predicted body weight) has traditionally been recommended to prevent hypoxemia and atelectasis in anesthetized patients.[6] There is, however, considerable evidence from experimental and observational studies that mechanical ventilation — in particular, high tidal volumes that cause alveolar overstretching — can initiate ventilator-associated lung injury[7] and contribute to extrapulmonary organ dysfunction through systemic release of inflammatory mediators.[8,9]

Lung-protective ventilation, which refers to the use of low tidal volumes and positive end-expiratory pressure (PEEP), and which may also include the use of recruitment maneuvers (periodic hyperinflation of the lungs),[10] has been shown to reduce mortality among patients with the acute respiratory distress syndrome[11] and is now considered best practice in the care of many critically ill patients.[12] Although this approach may be beneficial in a broader population,[13,14] some physicians have questioned the benefits of using lung-protective ventilation in the surgical setting,[15-18] especially since the use of high tidal volumes and no PEEP is still commonplace and less than 20% of patients receive protective ventilation in routine anesthetic practice.[19,20]

We conducted the Intraoperative Protective Ventilation (IMPROVE) trial to determine whether a multifaceted strategy of prophylactic lung-protective ventilation that combined low tidal volumes, PEEP, and recruitment maneuvers could improve outcomes after abdominal surgery, as compared with the standard practice of nonprotective mechanical ventilation.

Methods

Trial Design and Oversight

The IMPROVE trial was an investigator-initiated, multicenter, double-blind, stratified, parallel-group, clinical trial. Randomization was performed with the use of a computer-generated assignment sequence and a centralized telephone system. The study protocol and statistical analysis plan were approved for all centers by a central ethics committee (Comité de Protection des Personnes Sud-Est I, Saint-Etienne, France) according to French law. The protocol, including the statistical analysis plan, is available with the full text of this article at NEJM.org. An independent data and safety monitoring committee oversaw the study conduct and reviewed blinded safety data. The members of the steering committee (see the Supplementary Appendix, available at NEJM.org) vouch for the accuracy and completeness of the data and analyses and the fidelity of the study to the protocol. There was no industry support or involvement in the trial.

Patients were screened and underwent randomization between January 31, 2011, and August 10, 2012, at seven French university teaching hospitals. Written informed consent was obtained before randomization from each patient, on the day before surgery. Randomization was stratified according to study site and the planned use or nonuse of postoperative epidural analgesia, which is a factor that may influence outcomes.[21] Treatment assignments were concealed from patients, research staff, the statistician, and the data and safety monitoring committee. Although the staff members who collected data during surgery were aware of the group assignments, outcome assessors were unaware of these assignments throughout the study.

Patients

Patients were eligible for participation in the study if they were older than 40 years of age, were scheduled to undergo laparoscopic or nonlaparoscopic elective major abdominal surgery[1] with an expected duration of at least 2 hours, and had a preoperative risk index for pulmonary complications[5] of more than 2. The risk index uses risk classes that range from 1 to 5, with higher risk classes indicating a higher risk of postoperative pulmonary complications (see the

The New England Journal of Medicine is produced by NEJM Group, a division of the Massachusetts Medical Society.
Downloaded from nejm.org on October 14, 2026.

Supplementary Appendix). Patients were ineligible if they had received mechanical ventilation within the 2 weeks preceding surgery, had a body-mass index (the weight in kilograms divided by the square of the height in meters) of 35 or higher, had a history of respiratory failure or sepsis within the 2 weeks preceding surgery, had a requirement for intrathoracic or emergency surgery, or had a progressive neuromuscular illness.

INTERVENTIONS

Patients were assigned to receive volume-controlled mechanical ventilation according to one of two strategies: nonprotective ventilation with a tidal volume of 10 to 12 ml per kilogram of predicted body weight, with no PEEP and no recruitment maneuvers, as previously described[20] (the nonprotective-ventilation group), or lung-protective ventilation with a tidal volume of 6 to 8 ml per kilogram of predicted body weight, a PEEP of 6 to 8 cm of water, and recruitment maneuvers repeated every 30 minutes after tracheal intubation (the protective-ventilation group). Each recruitment maneuver consisted of applying a continuous positive airway pressure of 30 cm of water for 30 seconds. During anesthesia, a plateau pressure of no more than 30 cm of water was targeted in each group. All other ventilation procedures were identical in the two study groups (see the Supplementary Appendix).

The predicted body weight was calculated for each patient with the use of previously defined formulas.[11] For episodes of arterial desaturation (defined as a peripheral oxygen saturation of ≤92%), a transient increase in the fraction of inspired oxygen (F_{IO_2}) to 100% was permitted, and in patients assigned to nonprotective ventilation, the use of PEEP, recruitment maneuvers, or both was allowed, if required. Decisions about all other aspects of patient care during the intraoperative and postoperative periods, including general anesthesia, administration of fluids, use of prophylactic antibiotic agents, and postoperative pain management, were made by the attending physician according to the expertise of the staff at each center and routine clinical practice.

OUTCOMES

The primary outcome was a composite of major pulmonary and extrapulmonary complications occurring by day 7 after surgery. Major pulmonary complications were defined as pneumonia (defined according to standard criteria; see the Supplementary Appendix) or the need for invasive or noninvasive ventilation for acute respiratory failure. Major extrapulmonary complications were defined as sepsis, severe sepsis and septic shock (defined according to consensus criteria),[22] or death.

Secondary outcomes within the 30-day follow-up period were the incidence of pulmonary complications due to any cause, graded on a scale from 0 (no pulmonary complications) to 4 (the most severe complications)[23] (see the Supplementary Appendix); ventilation-related adverse events during surgery; postoperative gas exchange; unexpected need for admission to the intensive care unit (ICU); extrapulmonary complications; durations of ICU and hospital stays; and the rate of death from any cause 30 days after surgery. Pulmonary complications were analyzed separately; in particular, the need for invasive or noninvasive ventilation because of acute respiratory failure, the development of postoperative atelectasis, pneumonia, acute lung injury, and the acute respiratory distress syndrome, defined according to standard criteria (see the Supplementary Appendix). Extrapulmonary complications included the systemic inflammatory response syndrome (SIRS); sepsis; severe sepsis and septic shock; and surgical complications, including intraabdominal abscess, anastomotic leakage, and unplanned reoperation (all defined according to consensus criteria[22,24]).

STATISTICAL ANALYSIS

We calculated that a sample of 400 patients would provide 80% power to detect a relative difference of 50% in the primary outcome, at a two-sided alpha level of 0.05, assuming a 20% rate of postoperative complications in the nonprotective-ventilation group.[25] For safety reasons, an interim analysis was conducted after the enrollment of the first 200 patients, according to the a priori statistical analysis plan. The data and safety monitoring committee did not recommend discontinuation of the trial on the basis of that analysis, and 400 patients were therefore included. A total of 3 patients were excluded after randomization; surgery was stopped prematurely in 2 of the 3 patients because of extensive illness (duration of surgery, <2 hours), and 1 had undergone randomization in error (violation of exclusion criteria). An additional 3 patients were thus randomly assigned to a study group to obtain the full sample.

The New England Journal of Medicine is produced by NEJM Group, a division of the Massachusetts Medical Society.
Downloaded from nejm.org on October 14, 2026.

All analyses were conducted on data from the modified intention-to-treat population, which included all patients who underwent randomization except the three who were excluded (Fig. 1). An unadjusted chi-square test was used for the primary outcome analysis. Multiple logistic-regression analysis was used to identify relevant baseline covariates associated with the primary outcome, in addition to the stratification variables (use or nonuse of epidural analgesia and study center). Variables tested in the model were selected if the P value was less than 0.10 and if they were clinically relevant. Adjusted analyses were performed with the use of robust Poisson generalized-linear-model regression[26] and are presented as relative risks with 95% confidence intervals. A chi-square test (or Fisher's exact test, as appropriate) was used for secondary binary outcomes. The Hochberg procedure was used to adjust for multiple testing of components of the composite primary outcome.[27] Adjusted analyses were performed with the use of the same adjustment variables that were used in the robust Poisson regression analysis. Continuous variables were compared with the use of an unpaired t-test or the Mann–Whitney U test. Adjusted analyses were performed with the use




Figure 1. Assessment, Randomization, and Follow-up of Patients.

A total of 1803 patients awaiting abdominal surgery were assessed preoperatively for trial eligibility by the research staff. A total of 400 patients were included in the modified intention-to-treat analysis and were followed for 30 days after surgery. After randomization, 3 patients were excluded; 2 patients were excluded because surgery was stopped prematurely (duration, <2 hours), owing to extensive illness, and 1 had undergone randomization in error. An additional 3 patients were then enrolled in the study.

The New England Journal of Medicine is produced by NEJM Group, a division of the Massachusetts Medical Society.
Downloaded from nejm.org on October 14, 2026.

Table 1. Baseline Characteristics of the Patients.*

Characteristic	Nonprotective Ventilation (N=200)	Lung-Protective Ventilation (N=200)
Age — yr	63.4±10.0	61.6±11.0
Male sex — no. (%)	121 (60.5)	116 (58.0)
Height — cm	169.5±9.0	169.1±8.8
Body weight — kg		
Actual	71.3±13.9	71.4±14.2
Predicted†	63.8±9.9	63.3±9.7
Body-mass index‡		
Mean	24.7±3.8	24.8±3.8
25–35 — no. (%)	88 (44.0)	99 (49.5)
Preoperative risk index — no. (%)§		
Risk class 2	100 (50.0)	101 (50.5)
Risk class 3	94 (47.0)	93 (46.5)
Risk class 4 or 5	6 (3.0)	6 (3.0)
Coexisting condition — no. (%)¶		
Current smoking	50 (25.0)	51 (25.5)
Any alcohol intake	10 (5.0)	21 (10.5)
Not fully independent in activities of daily living	8 (4.0)	8 (4.0)
Chronic obstructive pulmonary disease	20 (10.0)	20 (10.0)
Loss of >10% of body weight in previous 6 mo	44 (22.0)	40 (20.0)
Long-term glucocorticoid use	4 (2.0)	7 (3.5)
Laparoscopic surgery — no. (%)	44 (22.0)	41 (20.5)
Type of surgery — no. (%)		
Pancreaticoduodenectomy	80 (40.0)	84 (42.0)
Liver resection	52 (26.0)	44 (22.0)
Gastrectomy	17 (8.5)	15 (7.5)
Colorectal resection	40 (20.0)	47 (23.5)
Other procedure	11 (5.5)	10 (5.0)
Diagnosis — no. (%)		
Cancer	164 (82.0)	155 (77.5)
Diagnosis other than cancer	36 (18.0)	45 (22.5)

* Plus–minus values are means ±SD. There were no significant differences between the two groups (P>0.05).

† The predicted body weight was calculated as follows: for men, 50+0.91(height in centimeters − 152.4); and for women, 45.5 + 0.91(height in centimeters − 152.4).[11]

‡ The body-mass index is the weight in kilograms divided by the square of the height in meters.

§ The preoperative risk index for pulmonary complications[5] uses risk classes that range from 1 to 5, with higher risk classes indicating a higher risk of postoperative complications. Patients with a risk class of 2 or more were eligible for participation in the study.

¶ All factors listed as coexisting conditions were included in the preoperative risk index as predictors of postoperative pulmonary complications.

of the same adjustment variables that were used in the linear-regression model. The time-to-event curves were calculated with the Kaplan–Meier method. Details regarding the handling of missing data are provided in the Supplementary Appendix.

All analyses were conducted with the use of Stata software, version 12 (StataCorp). A two-sided P value of less than 0.05 was considered to indicate statistical significance.

RESULTS

STUDY POPULATION

From January 2011 through August 2012, a total of 1803 patients awaiting abdominal surgery were assessed for trial eligibility. A total of 400 patients were included in the modified intention-to-treat analysis and were followed for 30 days after surgery (Fig. 1). One patient in the nonprotective-ventilation group received lung-protective ventilation but was included in the analysis for the group to which he was assigned. Data on the primary outcome were available for all patients. Baseline characteristics were similar between the two groups (Table 1). Open laparotomy, mainly for cancer resection, was performed in 156 patients (78.0%) in the nonprotective-ventilation group and in 159 (79.5%) in the protective-ventilation group (P=0.80).

INTRAOPERATIVE PROCEDURES

Table 2 shows the distribution of the main intraoperative procedures. Mean (±SD) tidal volumes were 11.1±1.1 ml per kilogram in the nonprotective-ventilation group, as compared with 6.4±0.8 ml per kilogram in the protective-ventilation group (P<0.001), and values remained within target ranges throughout the intraoperative period. In the protective-ventilation group, the median PEEP was 6 cm of water (interquartile range, 6 to 8), and the median number of recruitment maneuvers was 9 (interquartile range, 6 to 12); in the nonprotective-ventilation group, the value for each of these measures was 0 (interquartile range, 0 to 0) (Table 2). There were no significant between-group differences in type and duration of surgery, use or nonuse of epidural analgesia, blood loss, volume of fluids administered, and need for vasopressor administration. Five patients in the nonprotective-ventilation group required at least one intraoperative rescue therapy for arterial de-

The New England Journal of Medicine is produced by NEJM Group, a division of the Massachusetts Medical Society.
Downloaded from nejm.org on October 14, 2026.

Table 2. Intraoperative Procedures.*

Variable	Nonprotective Ventilation (N=200)	Lung-Protective Ventilation (N=200)	P Value
Tidal volume — ml	719.0±127.8	406.7±75.6	<0.001
Tidal volume — ml/kg of predicted body weight	11.1±1.1	6.4±0.8	<0.001
PEEP — cm of water			
Baseline			<0.001
Median	0	6	
Interquartile range	0–0	6–8	
End of surgery			<0.001
Median	0	6	
Interquartile range	0–0	6–8	
No. of recruitment maneuvers			<0.001
Median	0	9	
Interquartile range	0–0	6–12	
Peak pressure — cm of water			
Baseline	20.1±4.9	18.9±3.6	0.04
End of surgery	20.6±4.4	20.0±4.0	0.15
Plateau pressure — cm of water			
Baseline	16.1±4.3	15.2±3.0	0.02
End of surgery	16.6±3.5	15.2±2.6	<0.001
Respiratory system compliance — ml/cm of water			
Baseline	48.4±17.8	55.2±26.6	0.06
End of surgery	45.1±12.9	55.2±26.7	<0.001
F_{IO_2} — %	47.2±7.6	46.4±7.3	0.27
Volume of fluids administered — liters			
Crystalloid			0.47
Median	2.0	1.5	
Interquartile range	1.5–3.5	2.0–3.0	
Colloid			0.97
Median	0.5	0.5	
Interquartile range	0.25–1.0	0.50–1.0	
Duration of surgery — no./total no. (%)†			0.95
2–4 hr	76/192 (39.6)	75/195 (38.5)	
>4–6 hr	75/192 (39.1)	76/195 (39.0)	
>6 hr	41/192 (21.4)	44/195 (22.6)	
Duration of mechanical ventilation — min	344±127.9	319±139.4	0.84
Epidural analgesia — no. (%)	77 (38.5)	83 (41.5)	0.61

* Plus–minus values are means ±SD. Detailed data on intraoperative procedures are given in Table S2 in the Supplementary Appendix. F_{IO_2} denotes inspired oxygen fraction, and PEEP positive end-expiratory pressure.
† The duration of surgery was calculated as the time between skin incision and closure of the incision.

The New England Journal of Medicine is produced by NEJM Group, a division of the Massachusetts Medical Society.
Downloaded from nejm.org on October 14, 2026.

saturation (PEEP in one patient, recruitment maneuvers in two, and both in two), as compared with no patients in the protective-ventilation group (P=0.06).

OUTCOMES

Primary Outcome

Major pulmonary and extrapulmonary complications occurred within the first 7 days after surgery in 21 patients (10.5%) in the protective-ventilation group, as compared with 55 (27.5%) in the nonprotective-ventilation group (adjusted relative risk, 0.40; 95% confidence interval [CI], 0.24 to 0.68; P=0.001) (Table 3). The results of associated univariate and multivariate analyses are provided in Table S1 in the Supplementary Appendix.

Secondary Outcomes

One or more pulmonary complications developed within the first 7 days after surgery in 35 patients (17.5%) in the protective-ventilation group, as compared with 72 (36.0%) in the nonprotective-ventilation group (adjusted relative risk, 0.49; 95% CI, 0.32 to 0.74; P<0.001). More patients in the nonprotective-ventilation group than in the protective-ventilation group had major (grade ≥3) pulmonary complications (Table 3, and Tables S3 and S4 in the Supplementary Appendix) and major pulmonary and extrapulmonary complications during the 30 days after surgery (P<0.001 by the log-rank test) (Fig. 2). There were no relevant between-group differences in gas exchange after extubation and on day 1 after surgery (Table S5 in the Supplementary Appendix).

The proportion of patients who required postoperative ventilatory assistance (noninvasive ventilation or intubation) for acute respiratory failure was lower in the protective-ventilation group than in the nonprotective-ventilation group during the first 7 days after surgery (10 of 200 patients [5.0%], vs. 34 of 200 [17.0%]; adjusted relative risk, 0.29; 95% CI, 0.14 to 0.61; P=0.001), and the proportion was also lower with protective ventilation during the first 13 days after surgery (6.5% vs. 18.5%; adjusted relative risk, 0.36; 95% CI, 0.19 to 0.70; P=0.003) (Table 3). In addition, the cumulative 30-day probability of an event requiring intubation or noninvasive ventilation for postoperative acute respiratory failure was lower among patients who received lung-protective ventilation than among those who received nonprotective ventilation (P<0.001 by the log-rank test) (Fig. S1 in the Supplementary Appendix).

There was no significant difference between the protective-ventilation group and the nonprotective-ventilation group with respect to the proportion of patients who were unexpectedly admitted to the ICU during the 30-day period after surgery (11.0 and 12.5%, respectively; adjusted relative risk with protective ventilation, 0.88; 95% CI, 0.49 to 1.59; P=0.67), nor was there a significant between-group difference in the rate of adverse events (Table S3 in the Supplementary Appendix). Mortality at 30 days in the protective-ventilation group was similar to that in the nonprotective-ventilation group (3.0% and 3.5%, respectively; adjusted relative risk with protective ventilation, 1.13; 95% CI, 0.36 to 3.61; P=0.83). However, the median hospital stay was shorter in the protective-ventilation group than in the nonprotective-ventilation group (Table 3).

DISCUSSION

In this trial, intraoperative lung-protective mechanical ventilation, as compared with nonprotective ventilation, led to improved clinical outcomes and reduced health care utilization after abdominal surgery. The observed rate of postoperative complications in our study was slightly higher than predicted.[25] This was due, in part, to the exclusion of patients with a low risk of complications, as well as the large proportion of patients who underwent major abdominal procedures, which are associated with increased morbidity rates. Of the 400 patients enrolled, 19 had postoperative pneumonia and 47 had respiratory failure requiring intubation or noninvasive ventilation. These rates are consistent with previously reported rates of pulmonary complications[25,28] and mortality.[29] Our lung-protective ventilation strategy resulted in a 69% reduction in the number of patients requiring ventilatory support within the first 7 days after surgery.

Several hypotheses could explain some of the differences between the results of the present study and findings in other trials of lung-protective ventilation during high-risk surgery. Previous trials have included small numbers of patients, have focused on different (and not necessarily clinically relevant) outcomes,[17] and

The New England Journal of Medicine is produced by NEJM Group, a division of the Massachusetts Medical Society.
Downloaded from nejm.org on October 14, 2026.

Table 3. Results of Unadjusted and Adjusted Outcome Analyses.*

Variable	Nonprotective Ventilation (N=200)	Lung-Protective Ventilation (N=200)	Unadjusted Relative Risk or Between-Group Difference (95% CI)	P Value†	Adjusted Relative Risk or Between-Group Difference (95% CI)‡	P Value
Primary composite outcome — no. (%)						
Within 7 days§	55 (27.5)	21 (10.5)	0.38 (0.24–0.61)	<0.001	0.40 (0.24–0.68)	0.001
Within 30 days	58 (29.0)	25 (12.5)	0.43 (0.28–0.66)	<0.001	0.45 (0.28–0.73)	<0.001
Secondary outcomes — no. (%)						
Pulmonary complication within 7 days¶						
Grade 1 or 2	30 (15.0)	25 (12.5)	0.69 (0.42–1.13)	0.14	0.67 (0.39–1.16)	0.16
Grade ≥3	42 (21.0)	10 (5.0)	0.24 (0.12–0.46)	<0.001	0.23 (0.11–0.49)	<0.001
Atelectasis within 7 days‖	34 (17.0)	13 (6.5)	0.38 (0.21–0.70)	0.001	0.37 (0.19–0.73)	0.004
Pneumonia within 7 days	16 (8.0)	3 (1.5)	0.19 (0.05–0.63)	0.01	0.19 (0.05–0.66)	0.009
Acute lung injury or ARDS within 7 days	6 (3.0)	1 (0.5)	0.17 (0.02–1.37)	0.12	0.21 (0.02–1.71)	0.14
Need for ventilation within 7 days						
Invasive	7 (3.5)	2 (1.0)	0.29 (0.06–1.36)	0.51	0.40 (0.08–1.97)	0.26
Noninvasive	29 (14.5)	9 (4.5)	0.31 (0.15–0.64)	0.006	0.29 (0.13–0.65)	0.002
Extrapulmonary complication within 7 days						
SIRS	100 (50.0)	86 (43.0)	0.86 (0.70–1.06)	0.16	0.87 (0.65–1.17)	0.37
Sepsis	29 (14.5)	13 (6.5)	0.45 (0.24–0.84)	0.04	0.48 (0.25–0.93)	0.03
Severe sepsis or septic shock	9 (4.5)	8 (4.0)	0.89 (0.35–2.26)	0.80	1.48 (0.51–4.32)	0.47
Death within 30 days	7 (3.5)	6 (3.0)	0.86 (0.29–2.51)	0.80	1.13 (0.36–3.61)	0.83
Duration of stay in hospital and ICU — days						
Hospital				0.02		0.006
Median	13	11	−2.25 (−4.04 to −0.47)		−2.45 (−4.17 to −0.72)	
Interquartile range	8–20	8–15				
ICU				0.58		0.69
Median	7	6	−1.48 (−6.87 to 3.91)		−1.21 (−4.98 to 7.40)	
Interquartile range	4–9	4–8				

* All postoperative complications were defined according to consensus criteria (see the Supplementary Appendix). For additional data on postoperative outcomes, see Tables S3 and S4 in the Supplementary Appendix. ARDS denotes acute respiratory distress syndrome, CI confidence interval, ICU intensive care unit, and SIRS systemic inflammatory response syndrome. Relative risks are shown for outcome variables, and differences between groups are shown for the duration of stays in the hospital and ICU.

† Adjustment was performed for stratification variables (use or nonuse of epidural analgesia and study center), preoperative risk index for postoperative pulmonary complications, sex, duration of surgery, and need for blood transfusion (yes or no).

‡ The Hochberg procedure was used to adjust for multiple testing of components of the composite primary outcome.[27]


§ The primary outcome was a composite of major pulmonary complications (defined as pneumonia or need for invasive or noninvasive ventilation for acute respiratory failure) and extrapulmonary complications (defined as sepsis, septic shock, or death) within the first 7 days after surgery.

¶ Postoperative pulmonary complications were scored with the use of a graded scale[23] from 0 (no pulmonary complications) to 4 (the most severe complications) (see the Supplementary Appendix).

‖ Atelectasis was defined as opacification of the lung with shift of the mediastinum, hilum, or hemidiaphragm toward the affected area and compensatory overinflation in the adjacent, nonatelectatic lung.

have used either very low levels of PEEP or no PEEP.[15,16,18] One strength of the present trial is our use of a robust composite outcome that is highly pertinent to this high-risk surgical population.[5] Mechanical ventilation itself can induce an inflammatory response[30] and can synergize with the response induced by major surgery at both local and systemic levels. This amplification of

The New England Journal of Medicine is produced by NEJM Group, a division of the Massachusetts Medical Society.
Downloaded from nejm.org on October 14, 2026.



No. at Risk						
Nonprotective ventilation	200	182	163	145	142	142
Lung-protective ventilation	200	192	184	179	176	175

Figure 2. Kaplan–Meier Estimates of the Probability of the Composite Primary Outcome.

Data for the Kaplan–Meier estimates of the probability of the composite primary outcome of major pulmonary or extrapulmonary complications were censored at 30 days after surgery. Major pulmonary complications included pneumonia or the need for invasive or noninvasive ventilation for acute respiratory failure. Major extrapulmonary complications were sepsis, severe sepsis, septic shock, and death. P<0.001 by the log-rank test for the between-group difference in the probability of the primary outcome.

the inflammation cascade contributes to the subsequent development of lung injury[31] and systemic organ failure.[8,32]

The use of very low levels of PEEP in previous trials may have promoted the repeated opening and closing of small airways, leading to atelectasis, which can precipitate the development of pulmonary complications.[6,33] We used a multifaceted strategy of lung-protective ventilation that combined low tidal volumes, recruitment maneuvers to open collapsed alveoli, and moderate levels of PEEP to prevent further collapse.[34] Other strengths of the present trial include the methods used to minimize bias (blinded and centralized randomization, complete follow-up, and intention-to-treat analyses); the pragmatic nature of the trial protocol, with routine practice being maintained; and the enrollment of patients with characteristics similar to those of patients enrolled in other studies analyzing outcomes after major surgery.[29]

Our findings are consistent with the observation of transient arterial hypotension during recruitment maneuvers.[35] Consequently, recruitment maneuvers, in which hemodynamic effects are potentially influenced by the applied level of alveolar pressure,[36] should be used with caution in patients with hemodynamic instability.

There are several limitations to our study. The trial design did not include standardization of the administration of fluids. However, this limitation is unlikely to have affected our results, since the volume of fluids administered was similar in the two groups. The definition of nonprotective ventilation was arbitrary but is supported in the literature.[19,20] The trial protocol did not include standardization of requirements for noninvasive ventilation; however, it was recommended that the study centers follow clinical-practice guidelines,[37,38] and postoperative care was conducted by health care workers who were unaware of the study assignments. The utilization of noninvasive ventilation in our trial is close to that reported in earlier studies.[37] We therefore consider it unlikely that any imbalance in interventions affected our results.

In conclusion, our study provides evidence that a multifaceted strategy of prophylactic lung-protective ventilation during surgery, as compared with a practice of nonprotective mechanical ventilation, results in fewer postoperative complications and reduced health care utilization.

Dr. Futier reports receiving consulting fees from General Electric Medical Systems, lecture fees from Fresenius Kabi, and reimbursement of travel expenses from Fisher and Paykel Healthcare. Dr Constantin reports receiving consulting fees from Baxter, Fresenius Kabi, Dräger, and General Electric Medical Systems, payment for expert testimony from Baxter, Dräger, and Fresenius Kabi, lecture fees from General Electric Medical Systems, Dräger, Fresenius Kabi, Baxter, Hospal, Merck Sharp & Dohme, and LFB Biomedicaments, payment for the development of educational presentations from Dräger, General Electric Medical Systems, Baxter, and Fresenius Kabi, and reimbursement of travel expenses from Bird, Astute Medical, Astellas, Fresenius Kabi, Baxter, and Hospal. Dr. Paugam-Burtz reports receiving consulting fees from Fresenius Kabi, lecture fees and reimbursement of travel expenses from Astellas, and payment for the development of educational presentations from LFB Biomedicaments and Merck Sharp & Dohme. Dr. Allaouchiche reports receiving consulting fees from Fresenius Kabi and lecture fees from Novartis and Astellas. Dr. Leone reports receiving consulting fees from LFB Biomedicaments and lecture fees from Fresenius Kabi and Novartis. Dr. Jaber reports receiving consulting fees from Dräger France and Maquet France, lecture fees from Fisher and Paykel Healthcare, Abbott, and Philips, and reimbursement of travel expenses from Pfizer. No other potential conflict of interest relevant to this article was reported.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

We thank all the patients who participated in the study; the clinical and research staff at all the trial sites, without whose assistance the study would never have been completed; and Mervyn Singer for valuable advice during the preparation of the manuscript.

The New England Journal of Medicine is produced by NEJM Group, a division of the Massachusetts Medical Society.
Downloaded from nejm.org on October 14, 2026.

REFERENCES

1. Weiser TG, Regenbogen SE, Thompson KD, et al. An estimation of the global volume of surgery: a modelling strategy based on available data. Lancet 2008;372:139-44.
2. Khuri SF, Henderson WG, DePalma RG, et al. Determinants of long-term survival after major surgery and the adverse effect of postoperative complications. Ann Surg 2005;242:326-41.
3. Shander A, Fleisher LA, Barie PS, Bigtello LM, Sladen RN, Watson CB. Clinical and economic burden of postoperative pulmonary complications: patient safety summit on definition, risk-reducing interventions, and preventive strategies. Crit Care Med 2011;39:2163-72.
4. Arozullah AM, Daley J, Henderson WG, Khuri SF. Multifactorial risk index for predicting postoperative respiratory failure in men after major noncardiac surgery: the National Veterans Administration Surgical Quality Improvement Program. Ann Surg 2000;232:242-53.
5. Arozullah AM, Khuri SF, Henderson WG, Daley J. Development and validation of a multifactorial risk index for predicting postoperative pneumonia after major noncardiac surgery. Ann Intern Med 2001;135:847-57.
6. Bendixen HH, Hedley-Whyte J, Laver MB. Impaired oxygenation in surgical patients during general anesthesia with controlled ventilation: a concept of atelectasis. N Engl J Med 1963;269:991-6.
7. Serpa Neto A, Cardoso SO, Manetta JA, et al. Association between use of lung-protective ventilation with lower tidal volumes and clinical outcomes among patients without acute respiratory distress syndrome: a meta-analysis. JAMA 2012;308:1651-9.
8. Imai Y, Parodo J, Kajikawa O, et al. Injurious mechanical ventilation and end-organ epithelial cell apoptosis and organ dysfunction in an experimental model of acute respiratory distress syndrome. JAMA 2003;289:2104-12.
9. Lellouche F, Dionne S, Simard S, Bussieres J, Dagenais F. High tidal volumes in mechanically ventilated patients increase organ dysfunction after cardiac surgery. Anesthesiology 2012;116:1072-82.
10. Amato MB, Barbas CS, Medeiros DM, et al. Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome. N Engl J Med 1998;338:347-54.
11. The Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. N Engl J Med 2000;342:1301-8.
12. Putensen C, Theuerkauf N, Zinserling J, Wrigge H, Pelosi P. Meta-analysis: ventilation strategies and outcomes of the acute respiratory distress syndrome and acute lung injury. Ann Intern Med 2009;151:566-76. [Erratum, Ann Intern Med 2009;151:897.]
13. Schultz MJ, Haitsma JJ, Slutsky AS, Gajic O. What tidal volumes should be used in patients without acute lung injury? Anesthesiology 2007;106:1226-31.
14. Severgnini P, Selmo G, Lanza C, et al. Protective mechanical ventilation during general anesthesia for open abdominal surgery improves postoperative pulmonary function. Anesthesiology 2013 March 29 (Epub ahead of print).
15. Wrigge H, Uhlig U, Zinserling J, et al. The effects of different ventilatory settings on pulmonary and systemic inflammatory responses during major surgery. Anesth Analg 2004;98:775-81.
16. Wrigge H, Uhlig U, Baumgarten G, et al. Mechanical ventilation strategies and inflammatory responses to cardiac surgery: a prospective randomized clinical trial. Intensive Care Med 2005;31:1379-87.
17. Hong CM, Xu DZ, Lu Q, et al. Low tidal volume and high positive end-expiratory pressure mechanical ventilation results in increased inflammation and ventilator-associated lung injury in normal lungs. Anesth Analg 2010;110:1652-60.
18. Treschan TA, Kaisers W, Schaefer MS, et al. Ventilation with low tidal volumes during upper abdominal surgery does not improve postoperative lung function. Br J Anaesth 2012;109:263-71.
19. Jaber S, Coisel Y, Chanques G, et al. A multicentre observational study of intraoperative ventilatory management during general anaesthesia: tidal volumes and relation to body weight. Anaesthesia 2012;67:999-1008.
20. Hess DR, Kondili D, Burns E, Bittner EA, Schmidt UH. A 5-year observational study of lung-protective ventilation in the operating room: a single-center experience. J Crit Care 2013 January 29 (Epub ahead of print).
21. Pöpping DM, Elia N, Marret E, Remy C, Tramèr MR. Protective effects of epidural analgesia on pulmonary complications after abdominal and thoracic surgery: a meta-analysis. Arch Surg 2008;143:990-9.
22. Bone RC, Sibbald WJ, Sprung CL. The ACCP-SCCM consensus conference on sepsis and organ failure. Chest 1992;101:1481-3.
23. Hulzebos EH, Helders PJ, Favié NJ, et al. Preoperative intensive inspiratory muscle training to prevent postoperative pulmonary complications in high-risk patients undergoing CABG surgery: a randomized clinical trial. JAMA 2006;296:1851-7.
24. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. Ann Surg 2004;240:205-13.
25. Thompson JS, Baxter BT, Allison JG, Johnson FE, Lee KK, Park WY. Temporal patterns of postoperative complications. Arch Surg 2003;138:596-602.
26. Zou G. A modified Poisson regression approach to prospective studies with binary data. Am J Epidemiol 2004;159:702-6.
27. Hochberg Y. A sharper Bonferroni procedure for multiple tests of significance. Biometrika 1988;75:800-2.
28. Lawrence VA, Dhanda R, Hilsenbeck SG, Page CP. Risk of pulmonary complications after elective abdominal surgery. Chest 1996;110:744-50.
29. Pearse RM, Moreno RP, Bauer P, et al. Mortality after surgery in Europe: a 7 day cohort study. Lancet 2012;380:1059-65.
30. Ranieri VM, Suter PM, Tortorella C, et al. Effect of mechanical ventilation on inflammatory mediators in patients with acute respiratory distress syndrome: a randomized controlled trial. JAMA 1999;282:54-61.
31. Mascia L, Pasero D, Slutsky AS, et al. Effect of a lung protective strategy for organ donors on eligibility and availability of lungs for transplantation: a randomized controlled trial. JAMA 2010;304:2620-7.
32. Bouadma L, Dreyfuss D, Ricard JD, Martet G, Saumon G. Mechanical ventilation and hemorrhagic shock-resuscitation interact to increase inflammatory cytokine release in rats. Crit Care Med 2007;35:2601-6.
33. Duggan M, Kavanagh BP. Pulmonary atelectasis: a pathogenic perioperative entity. Anesthesiology 2005;102:838-54.
34. Futier E, Constantin JM, Pelosi P, et al. Noninvasive ventilation and alveolar recruitment maneuver improve respiratory function during and after intubation of morbidly obese patients: a randomized controlled study. Anesthesiology 2011;114:1354-63.
35. Fan E, Wilcox ME, Brower RG, et al. Recruitment maneuvers for acute lung injury: a systematic review. Am J Respir Crit Care Med 2008;178:1156-63.
36. Lim SC, Adams AB, Simonson DA, et al. Transient hemodynamic effects of recruitment maneuvers in three experimental models of acute lung injury. Crit Care Med 2004;32:2378-84.
37. Jaber S, Chanques G, Jung B. Postoperative noninvasive ventilation. Anesthesiology 2010;112:453-61.
38. Keenan SP, Sinuff T, Burns KE, et al. Clinical practice guidelines for the use of noninvasive positive-pressure ventilation and noninvasive continuous positive airway pressure in the acute care setting. CMAJ 2011;183(3):E195-E214.

Copyright © 2013 Massachusetts Medical Society.

The New England Journal of Medicine is produced by NEJM Group, a division of the Massachusetts Medical Society.
Downloaded from nejm.org on October 14, 2026.